

A Note on Wess-Zumino Terms and Discrete Symmetries

Silas R. Beane

Department of Physics, University of Maryland
College Park, MD 20742-4111

sbeane@pion.umd.edu

ABSTRACT

Sigma models in which the integer coefficient of the Wess-Zumino term vanishes are easy to construct. This is the case if all flavor symmetries are vectorlike. We show that there is a subset of $SU(N) \times SU(N)$ vectorlike sigma models in which the Wess-Zumino term vanishes for reasons of symmetry as well. However, there is no chiral sigma model in which the Wess-Zumino term vanishes for reasons of symmetry. This can be understood in the sigma model basis as a consequence of an index theorem for the axialvector coupling matrix. We prove this index theorem directly from the $SU(N) \times SU(N)$ algebra.

PACS: 11.30.Rd; 12.38.Aw; 12.90.+b; 11.30.Er

1. Introduction

Any microscopic theory which exhibits the pattern of spontaneous symmetry breaking $SU(N) \times SU(N) \rightarrow SU(N)$ is described at low energies by an effective theory of $N^2 - 1$ Goldstone bosons living on the coset space $SU(N) \times SU(N)/SU(N)$. Well known technology tells how to build the most general lagrangian involving the Goldstone bosons [1]. Introduce a field U that transforms linearly with respect to $SU(N)_L \times SU(N)_R$: $U \rightarrow LUR^\dagger$, where L, R is an element of $SU(N)_{L,R}$. A convenient parametrization of U is

$$U = \exp \frac{2i\pi_a T_a}{F_\pi} \quad (1)$$

where the T_a are $SU(N)$ generators normalized such that $\text{Tr}(T_a T_b) = \delta_{ab}/2$ and π_a is the canonical Goldstone boson field with decay constant F_π . In this basis the Goldstone bosons transform nonlinearly with respect to $SU(N)_L \times SU(N)_R$. Of the manifestly invariant terms in the lagrangian, the unique term with the fewest derivatives is

$$\frac{1}{4} F_\pi^2 \text{Tr} \partial_\mu U \partial^\mu U^\dagger. \quad (2)$$

Consider the discrete symmetries of this operator [2]. It is invariant with respect to charge conjugation, $U \leftrightarrow U^T$, and parity, $U_{\vec{x}} \leftrightarrow U_{-\vec{x}}^\dagger$. However, the operator is separately invariant with respect to $U \leftrightarrow U^\dagger$, or $\pi \rightarrow -\pi$, which counts modulo two the number of bosons. Therefore this operator, as well as all other manifestly invariant terms, has a discrete symmetry, unrelated to P, C and T , which forbids interactions of odd numbers of Goldstone bosons. The absence of such a discrete symmetry in realistic underlying theories like QCD can be used to motivate the existence of other terms which involve interactions of odd numbers of Goldstone modes [2].

Although these so-called Wess-Zumino terms are not manifestly invariant, they transform to a total derivative, which leaves the action invariant [3]. They are the unique terms which transform to a total derivative [4]. General topological reasoning reveals that the coefficient of such operators must be an integer [2]. This argument follows from the topology of the coset space and is therefore independent of details of the microscopic theory. The contribution of the Wess-Zumino term to the action can be written as

$$n\Gamma_{WZ} \quad (3)$$

where n is an integer [2]. Note that Γ_{WZ} is subject to usual operator rules of thumb. For instance, it vanishes only if there is a symmetry —such as $U \leftrightarrow U^\dagger$ — which forbids it. For this note it is relevant to know that Γ_{WZ} contains interactions of odd numbers of Goldstone bosons and the gauged Wess-Zumino term gives rise to the process $\pi_0 \rightarrow \gamma\gamma$.

$$\sum_i \text{---} \pi_0 \text{---} \triangle \text{---} \gamma \gamma \sim \text{Tr}(\hat{g}_A)$$

Figure 1: In a sigma model with arbitrary fermion content the amplitude for $\pi_0 \rightarrow \gamma\gamma$ is proportional to $\text{Tr}(\hat{g}_A)$.

The value of the integer n is found by matching to the microscopic theory. If $SU(N) \times SU(N)$ is chiral in the underlying gauge theory then there are nonvanishing flavor anomalies. Here all axial symmetries are by assumption spontaneously broken and therefore there is no 't Hooft matching of anomalies between the microscopic theory and the low-energy theory [5]. However, anomalous Ward identities must be satisfied. The Wess-Zumino term fulfills this role. If $SU(N) \times SU(N)$ is vectorlike in the underlying theory then there are no flavor anomalies since a fermion with a mass term allowed by a symmetry cannot contribute to an anomaly for that symmetry. We then expect a vanishing integer in front of the Wess-Zumino term. One might worry that the vectorlike scenario is in contradiction with the Vafa-Witten theorem [6], which constrains the breaking of vectorlike symmetries. This is not the case. A gauge theory with an $SU(N) \times SU(N)$ vectorlike flavor symmetry does not satisfy the Vafa-Witten positivity conditions. As we will see below, there are either fundamental scalars or nonrenormalizable operators in such a theory.

In QCD $SU(N) \times SU(N)$ is chiral and there are no discrete symmetries beyond P , C and T . Therefore, QCD is an example of an underlying theory in which neither n nor Γ_{WZ} vanish. To our knowledge, other possibilities have not been considered in the literature. In this note we construct models in which n vanishes and Γ_{WZ} does not vanish and in which n and Γ_{WZ} vanish. Not surprisingly the fermion content of these models is vectorlike with respect to $SU(N) \times SU(N)$. We are unable to find a more interesting model in which n is nonzero and Γ_{WZ} vanishes. We will show that such a chiral model would be inconsistent.

It is convenient to work with sigma models. We focus entirely on $SU(N) \times SU(N)$ sigma models with fermions in the fundamental representation of $SU(N)$. Given any sigma model, we can always construct an asymptotically free gauge theory with precisely the same symmetry structure and anomaly content. In a sigma model fermions transform linearly with respect to $SU(N) \times SU(N)$ whereas Wess-Zumino terms appear in the nonlinear realization. Wess-Zumino terms can be derived from the sigma models by taking into account the effect of a change of basis on the path integral measure [7]. The effect of the Wess-Zumino term in the sigma model basis can be obtained by computing triangle

graphs with fermions on the internal lines. For instance, one can compute the amplitude for $\pi_0 \rightarrow \gamma\gamma$ [8]. Generally, in a sigma model with Goldstone bosons and any number of fermions, this process is proportional to $Tr(\hat{g}_A)$ where $\hat{g}_A$ is an axialvector coupling matrix (see Fig. 1). In order to reproduce the same physics in the two bases, it follows from Eq. (3) that

$$Tr(\hat{g}_A) = n \quad (4)$$

where n is the coefficient of the Wess-Zumino term. We will derive this index theorem for $\hat{g}_A$ directly from the $SU(N) \times SU(N)$ algebra, and we will show that as a consequence of this index theorem there is no chiral model with a discrete symmetry which implies a vanishing Wess-Zumino term.

In section 2 we investigate several chiral sigma models; in particular, we find the axialvector coupling matrices and search for points of enhanced symmetry. We investigate a vectorlike sigma model in section 3. In section 4, we generalize our results and state an index theorem for the axialvector coupling matrix. We show, as consequence of the index theorem, that there is no chiral sigma model with a discrete symmetry which implies a vanishing Wess-Zumino term. We then prove the theorem and discuss other possible applications and extensions. In section 5 we summarize.

2. The Chiral Models

2.1 Model I

The sigma models we will consider all contain a scalar field, Σ , transforming in the $(\bar{N}, N)$ representation of $SU(N) \times SU(N)$ together with various combinations of quark matter transforming in the $(1, N)$ and $(N, 1)$ representations¹. In this linear basis the discrete symmetry we are searching for involves the transformation $\Sigma \leftrightarrow \Sigma^\dagger$ together with a transformation of the quark matter. Consider first the ordinary sigma model. This chiral model reflects the symmetries and anomaly structure of N -flavor QCD. Its matter content consists of quarks and mesons transforming as:

	$SU(N)_L$	$SU(N)_R$
Q_L	N	1
Q_R	1	N
Σ	$\bar{\mathbf{N}}$	N

¹ Models similar to those discussed below have been investigated in Ref. 9.

with respect to $SU(N) \times SU(N)$. In our convention, $\mathcal{Q}_L = \frac{1}{2}(1+\gamma_5)\mathcal{Q}$ and $\mathcal{Q}_R = \frac{1}{2}(1-\gamma_5)\mathcal{Q}$. The invariant interaction is

$$\bar{\mathcal{Q}}_L \Sigma \mathcal{Q}_R + p.c. \quad (5)$$

It is clear that there is no additional discrete symmetry beyond QCD symmetries. The axialvector coupling of this model is easy to find by choosing $\Sigma = F_\pi U$ where U is defined in Eq. (1). This choice allows us to ignore details of the symmetry breaking mechanism. We find $g_A = 1$ and so $\pi_0 \rightarrow \gamma\gamma$ is nonvanishing and easily calculable from triangle graphs [10] for any number of flavors. Therefore, the Wess-Zumino term computed from this model by a change of basis is nonvanishing with $n = 1$ [7]. The correct QCD result follows from assigning each quark a color quantum number. In that case we have $n = N_c g_A = N_c$. Nonrenormalizable operators, such as

$$\bar{\mathcal{Q}}_L \Sigma i \not{\partial} \Sigma^\dagger \mathcal{Q}_L + p.c. \quad (6)$$

shift the value of g_A . However, they do not contribute to $\pi_0 \rightarrow \gamma\gamma$ in the chiral limit. Therefore they have nothing to do with the Wess-Zumino term in the transformed basis. It is sufficient to keep contributions to g_A from renormalizable operators. The significance of the renormalizability constraint will become clear in Section 4.

2.2 Model II

Adding more matter guarantees extra discrete symmetries for special values of couplings. Hence, consider another chiral sigma model with more matter, transforming as:

	$SU(N)_L$	$SU(N)_R$
$\mathcal{Q}_L, \tilde{\mathcal{Q}}_L$	N	1
$\mathcal{Q}_R, \tilde{\mathcal{Q}}_R$	1	N
Σ	$\bar{\mathbf{N}}$	N

with respect to $SU(N) \times SU(N)$. The invariant interaction is

$$a(\bar{\mathcal{Q}}_L \Sigma \mathcal{Q}_R + p.c.) + b(\tilde{\bar{\mathcal{Q}}}_L \Sigma \tilde{\mathcal{Q}}_R + p.c.) + c(\bar{\mathcal{Q}}_L \Sigma \tilde{\mathcal{Q}}_R + \tilde{\bar{\mathcal{Q}}}_L \Sigma \mathcal{Q}_R + p.c.). \quad (7)$$

With $a = b$ this model has a discrete symmetry with respect to interchange of $\mathcal{Q}_L$ and $\tilde{\mathcal{Q}}_R$, $\mathcal{Q}_R$ and $\tilde{\mathcal{Q}}_L$, and Σ and $\Sigma^\dagger$. This symmetry is a generalization of parity and leads to no new multiplicatively conserved quantum number. In this model we find,

$$\hat{g}_A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad (8)$$

for all values of the couplings a , b and c . Therefore $Tr(\hat{g}_A) = 2$, and $\pi_0 \rightarrow \gamma\gamma$ is nonvanishing and twice the value obtained in the ordinary sigma model. Hence the Wess-Zumino term has coefficient $n = 2$.

2.3 Model III

The chiral models considered above have no nontrivial mixing because there are no invariant quark bilinears. Consider therefore a chiral sigma model with yet more matter, transforming as:

	$SU(N)_L$	$SU(N)_R$
$Q_L, \tilde{Q}_L, P_R$	N	1
$Q_R, \tilde{Q}_R, P_L$	1	N
Σ	$\bar{\mathbf{N}}$	N

with respect to $SU(N) \times SU(N)$. The invariant interaction is:

$$\begin{aligned}
& a(\bar{Q}_L \Sigma Q_R + p.c.) + b(\tilde{\bar{Q}}_L \Sigma \tilde{Q}_R + p.c.) + c(\bar{Q}_L \Sigma \tilde{Q}_R + \tilde{\bar{Q}}_L \Sigma Q_R + p.c.) \\
& d(\bar{P}_R \Sigma P_L + p.c.) + M_0(\bar{P}_R Q_L + \bar{Q}_L P_R + p.c.) + \tilde{M}_0(\bar{P}_R \tilde{Q}_L + \tilde{\bar{Q}}_L P_R + p.c.).
\end{aligned} \tag{9}$$

Note the invariant bilinears. In the symmetric phase, two of the quarks are massive and degenerate and one is massless. Here there is a discrete symmetry with respect to interchange of Q and P and Σ and $\Sigma^\dagger$ when $a = d$, $b = c = \tilde{M}_0 = 0$, and with respect to interchange of $\tilde{Q}$ and P and Σ and $\Sigma^\dagger$ when $b = d$, $a = c = M_0 = 0$. However these choices of parameters correspond to decoupling $\tilde{Q}$ and Q , respectively, which yields the simplest vectorlike model for the remaining matter. We will study this model in the next section. In this model we find

$$\hat{g}_A = \begin{pmatrix} \cos^2 \gamma \cos 2\beta + \sin^2 \gamma & \cos \gamma \sin \gamma (\cos 2\beta - 1) & \cos \gamma \sin 2\beta \\ \cos \gamma \sin \gamma (\cos 2\beta - 1) & \sin^2 \gamma \cos 2\beta + \cos^2 \gamma & -\sin \gamma \sin 2\beta \\ \cos \gamma \sin 2\beta & -\sin \gamma \sin 2\beta & -\cos 2\beta \end{pmatrix}, \tag{10}$$

where the mixing angles are related to the coupling parameters. Note that $(\hat{g}_A)^2 = \mathbf{1}$. This is a statement of the Adler-Weisberger sum rule. If for special values of the couplings there is a discrete symmetry which implies a multiplicatively conserved quantum number and involves $\pi \rightarrow -\pi$, then the diagonal elements of $\hat{g}_A$ must vanish for the corresponding values of the mixing angles. But note that $Tr(\hat{g}_A) = 1$, independent of the values of the mixing angles. This is of course consistent with the index theorem. This model therefore has no relevant discrete symmetry, as deduced directly from the lagrangian. The amplitude

for $\pi_0 \rightarrow \gamma\gamma$ is the same as in the ordinary sigma model (Model I). We will see below that generally there is no chiral model with the extra discrete symmetry. But first we will consider the simplest vectorlike model.

3. The Vectorlike Model

The matter content of the vectorlike model consists of quarks and mesons transforming as:

	$SU(N)_L$	$SU(N)_R$
$\mathcal{Q}_L, \mathcal{P}_R$	N	1
$\mathcal{Q}_R, \mathcal{P}_L$	1	N
Σ	$\bar{\mathbf{N}}$	N

with respect to $SU(N) \times SU(N)$. Note that there are equal numbers of left- and right-handed Weyl fermions assigned to each charge in a vectorlike model. This guarantees that all quarks have invariant masses. The invariant interaction is

$$a(\bar{\mathcal{Q}}_L \Sigma \mathcal{Q}_R + p.c.) + b(\bar{\mathcal{P}}_R \Sigma \mathcal{P}_L + p.c.) + M_0(\bar{\mathcal{Q}}_L \mathcal{P}_R + \bar{\mathcal{Q}}_R \mathcal{P}_L + p.c.). \quad (11)$$

All fermions are massive. The free fermion theory is $U(2N) \times U(2N)$ invariant. The invariant mass term breaks this to $U(2N)$, and the Yukawa interactions further break this to $SU(N) \times SU(N) \times U(1)$. Therefore, unlike the ordinary chiral sigma model where a microscopic theory in which the fermions have only gauge interactions can always be constructed, here a microscopic theory must have additional nongauge interactions. It must therefore have fundamental scalars or nonrenormalizable interactions. In neither case does the Vafa-Witten theorem constrain the pattern of symmetry breaking [6]. Vectorlike $SU(N) \times SU(N)$ sigma models are therefore worthy of study.

Here we have an additional discrete symmetry if $a = b$. If $a = b$ there is a $\mathbf{Z}_2$ symmetry corresponding to interchange of $\mathcal{Q}$ and $\mathcal{P}$ and Σ and $\Sigma^\dagger$. This symmetry commutes with parity and so we can assign multiplicatively conserved charges to each physical state:

	P	$\mathbf{Z}_2$
ψ_+	1	1
ψ_-	1	-1
π	-1	-1

Here $\psi_{\pm}$ denote the quarks in the diagonal basis. This is precisely the discrete symmetry which rules out self interactions of odd numbers of Goldstone bosons. In this model we find

$$\hat{g}_A = \begin{pmatrix} -\cos 2\phi & -\sin 2\phi \\ -\sin 2\phi & \cos 2\phi \end{pmatrix}, \quad (12)$$

where $\cot 2\phi = (a - b)F_{\pi}/2M_0$. Again $(\hat{g}_A)^2 = \mathbf{1}$, a statement of the Adler-Weisberger sum rule. Since $Tr(\hat{g}_A) = 0$, $\pi_0 \rightarrow \gamma\gamma$ vanishes for all values of a and b , as expected in a vectorlike model. In the transformation to the nonlinear basis, the nontrivial Jacobians due to $\mathcal{Q}$ and $\mathcal{P}$ cancel. Note that if $a = b$,

$$\hat{g}_A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \quad (13)$$

The diagonal elements of the axialvector coupling matrix vanish because they do not respect the $\mathbf{Z}_2$ symmetry. That is, $g_{\pi\psi_-\psi_-} = g_{\pi\psi_+\psi_+} = 0$. The Adler-Weisberger sum rule then predicts the off-diagonal axialvector couplings. At this point of enhanced symmetry, all triangle graphs vanish (and not just the sum). Correspondingly, the Wess-Zumino operator in the nonlinear basis vanishes (and not just the integer coefficient). Clearly this is not of great practical interest.

This sigma model provides a field theoretic realization of Weinberg's theory of the axialvector coupling of the constituent quark [11]. The vectorlike nature of the model and the point of enhanced $\mathbf{Z}_2$ symmetry are in precise correspondence with algebraic sum rules for Goldstone boson scattering, which are well satisfied in nature [12]. This correspondence is a profound puzzle.

4. An Index Theorem

The results of the sigma models considered above can be summarized by the index theorem:

$$Tr(\hat{g}_A) = n_L - n_R \quad (14)$$

where, by convention, n_L, n_R is the number of left-handed Weyl fermions with $SU(N) \times SU(N)$ charge $(N, 1), (1, N)$. In a chiral sigma model $n_L \neq n_R$ and in a vectorlike model $n_L = n_R$, by definition. If $\pi \rightarrow -\pi$ with respect to a discrete symmetry which implies a multiplicatively conserved charge, then the diagonal elements of $\hat{g}_A$ transform like π . Therefore in a model with such a symmetry, $[\hat{g}_A]_{\alpha\alpha} = 0 \forall \alpha$, and clearly $Tr(\hat{g}_A) = 0$. Such a model is vectorlike. Therefore, there can be no chiral sigma model with such a discrete symmetry.

It is important to know precisely what enters Eq. (14). Here we give a proof of Eq. (14) that is purely a consequence of the $SU(N) \times SU(N)$ representation theory. We emphasize that the index theorem in the linear basis has nothing to do with topology. Assume that we are in a helicity conserving Lorentz frame and that the quarks, for each helicity λ , fall into a reducible representation of $SU(N) \times SU(N)$ consisting of a sum of any number of $(1, N)$ and $(N, 1)$ representations. For a given helicity, we can express the diagonal quark states as

$$|\psi\rangle_\alpha^\lambda = \sum_{j=1}^{n_L} U_{\alpha j} |N, 1\rangle_j + \sum_{\bar{k}=1}^{n_R} V_{\alpha \bar{k}} |1, N\rangle_{\bar{k}} \quad (15)$$

where $\hat{U}$ and $\hat{V}$ are mixing matrices and α ranges from 1 to $n_L + n_R$. The parity operator in the helicity conserving frame acts as $P|\psi\rangle_\alpha^\lambda = |\psi\rangle_\alpha^{-\lambda}$ and as $P|\alpha, \beta\rangle_i = |\beta, \alpha\rangle_i$ and $P|\alpha, \beta\rangle_{\bar{k}} = |\beta, \alpha\rangle_{\bar{k}}$ on states of definite $SU(N) \times SU(N)$. The $SU(N) \times SU(N)$ algebra can be expressed as

$$[\mathcal{Q}_a^5, \mathcal{Q}_b^5] = if_{abc} T_c \quad [T_a, \mathcal{Q}_b^5] = if_{abc} \mathcal{Q}_c^5 \quad [T_a, T_b] = if_{abc} T_c \quad (16)$$

where $\mathcal{Q}_b^5$ is the axialvector charge operator and the T_c are $SU(N)$ generators. The action of the axialvector charge operator on the states of definite $SU(N) \times SU(N)$ is

$$\mathcal{Q}_a^5 |1, N\rangle = -T_a |1, N\rangle \quad \mathcal{Q}_a^5 |N, 1\rangle = T_a |N, 1\rangle. \quad (17)$$

This sign convention is in agreement with Eq. (14). The quark axialvector coupling matrix is then defined by

$$\lambda \langle \psi | \mathcal{Q}_a^5 | \psi \rangle_\beta^\lambda = [\hat{g}_A^\lambda]_{\alpha\beta} T_a. \quad (18)$$

The results of the models of the previous sections for $\hat{g}_A$ are easily recovered by choosing values of n_L and n_R and a representation of the $\hat{U}$ and $\hat{V}$ mixing matrices. For instance, model I corresponds to $n_L = 1$, $n_R = 0$ and $U = 1$. Model III corresponds to $n_L = 2$, $n_R = 1$, and $\hat{U}$ and $\hat{V}$ a combination of Euler angles. The vectorlike model corresponds to $n_L = n_R = 1$, $U_1 = V_2 = \sin \phi$ and $V_1 = -U_2 = \cos \phi$. Therefore, all that is of interest in the sigma models is a consequence of the $SU(N) \times SU(N)$ representation theory. We are now in a position to prove the index theorem.

Proof:

Orthonormality of the $|\psi\rangle_\alpha$ and the definition of $\hat{g}_A$ give, respectively:

$$\sum_{j=1}^{n_L} U_{\alpha j}^* U_{\beta j} + \sum_{\bar{k}=1}^{n_R} V_{\alpha \bar{k}}^* V_{\beta \bar{k}} = \delta_{\alpha\beta} \quad (19a)$$

$$\sum_{j=1}^{n_L} U_{\alpha j}^* U_{\beta j} - \sum_{\bar{k}=1}^{n_R} V_{\alpha \bar{k}}^* V_{\beta \bar{k}} = [\hat{g}_A^\lambda]_{\alpha\beta}. \quad (19b)$$

It is easy to check that the generalized Adler-Weisberger sum rule, $[\hat{g}_A^\lambda]_{\alpha\beta} [\hat{g}_A^\lambda]_{\beta\gamma} = \delta_{\alpha\gamma}$, is satisfied. Since $Tr(\delta_{\alpha\beta}) = n_L + n_R$, Eq. (19a) and the independence of left and right give:

$$Tr \sum_{j=1}^{n_L} U_{\alpha j}^* U_{\beta j} = n_L \quad Tr \sum_{\bar{k}=1}^{n_R} V_{\alpha \bar{k}}^* V_{\beta \bar{k}} = n_R. \quad (20)$$

It then follows directly from Eq. (19b) that

$$Tr(\hat{g}_A^{\pm\lambda}) = \pm(n_L - n_R) \quad (21)$$

with $+\lambda$ identified with left-handed helicity and $-\lambda$ identified with right-handed helicity, which is the desired relation. *Q.E.D.*

It might seem puzzling that the index theorem follows purely from $SU(N) \times SU(N)$ while the sigma model results rely on renormalizability. There is no paradox; the full $SU(N) \times SU(N)$ symmetry in helicity conserving frames implies a constraint on the asymptotic behaviour of Goldstone boson scattering amplitudes in the broken phase [13]. Evidently renormalizability, which is of course also a statement about high energy behavior, yields an equivalent constraint in this context [11].

Does the index theorem generalize to higher representations of $SU(N) \times SU(N)$? For instance, in $N = 2$ QCD, baryons fall into reducible representations which are sums of any number of $(1, 2)$, $(2, 1)$ and $(1, 4)$, $(4, 1)$, $(3, 2)$ and $(2, 3)$ representations. The four-dimensional representations contain the isospin $3/2$ baryons. It is not clear to the author whether an extension of the index theorem to include these representations is possible or even sensible.

5. Summary

In the nonlinear realization of $SU(N) \times SU(N)/SU(N)$, manifestly invariant operators in the lagrangian have a discrete symmetry which rules out interactions of odd numbers of Goldstone bosons [2]. The fact that this discrete symmetry is not a symmetry of realistic theories like QCD motivates the existence of additional operators which break this discrete symmetry [3]. These Wess-Zumino operators are not manifestly invariant and yet they transform to a total derivative and therefore leave the action invariant. The coefficient of these operators is an integer, which is determined in practice by matching to the underlying theory. Generally in underlying theories like QCD in which $SU(N) \times SU(N)$ is chiral, this integer is nonvanishing. In this letter we have searched for a chiral model in which the

integer of the Wess-Zumino term is nonvanishing and yet has a discrete symmetry, $\pi \rightarrow -\pi$, which implies that the Wess-Zumino term must vanish.

We found it convenient to work with sigma models in which the fermion fields transform linearly with respect to $SU(N) \times SU(N)$. Wess-Zumino terms can be derived in these models by a transformation to a nonlinear basis [7]. In the linear models the effects of anomalies are contained in triangle graphs [8]. The fact that the coefficient of the Wess-Zumino term is an integer has an analogous statement in the linear basis. In the linear basis, the trace of the axialvector coupling matrix, which determines the sum of all triangle graphs, must be an integer. We proved this index theorem directly from the $SU(N) \times SU(N)$ algebra. In chiral models the integer is nonvanishing and in vectorlike models the integer vanishes. If a given sigma model has a discrete symmetry which implies a multiplicatively conserved charge and if $\pi \rightarrow -\pi$ with respect to this discrete symmetry, then each diagonal element of the axialvector coupling matrix must vanish. In such a case the trace vanishes. Therefore a model with this discrete symmetry must be vectorlike with respect to $SU(N) \times SU(N)$. We conclude that there is no chiral model with a discrete symmetry which implies a vanishing Wess-Zumino term.

This work was supported by the U.S. Department of Energy grant DE-FG02-93ER-40762. I thank Markus Luty for valuable conversations.

References

1. C.G. Callan, S. Coleman, J. Wess and B. Zumino, Phys. Rev. 177 (1969) 2247 .
2. E. Witten, Nucl. Phys. B 223 (1983) 422 .
3. J. Wess and B. Zumino, Phys. Lett. B 37 (1971) 95 .
4. E. D'Hoker and S. Weinberg, Phys. Rev. D 50 (1994) 6050 , [hep-ph/9409402](#).
5. G. 't Hooft, in *Recent Developments in Gauge Theories*, Proc. of the NATO ASI, Cargèse 1979, ed. G.'t Hooft *et al* (Plenum, 1980).
6. C. Vafa and E. Witten, Nucl. Phys. B 234 (1984) 173 .
7. A. Manohar and G. Moore, Nucl. Phys. B 243 (1984) 55 .
8. J. Steinberger, Phys. Rev. 76 (1949) 1180 .
9. C. deTar and T. Kunihiro, Phys. Rev. D 39 (1989) 2805 ; Y. Nemoto *et al*, [hep-ph/9710445](#).
10. C. Itzykson and J-B. Zuber, *Quantum Field Theory* (McGraw-Hill, 1980).
11. S. Weinberg, Phys. Rev. Lett. 65 (1990) 1177 ; *ibid*, 1181.
12. S.R. Beane, Ann. Phys. 263 (1998) 214 , [hep-ph/9706246](#).
13. S. Weinberg, Phys. Rev. 177 (1969) 2604 .